

On Phi-Pseudo-Monotonicity and Approximation-Solvability of Nonlinear Equations

RAM VERMA AND LOKENATH DEBNATH

Department of Mathematics, University of Central Florida, Orlando, Florida 32816

Submitted by E. W. Cheney

Received April 13, 1992

We generalize the results on the approximation-solvability of nonlinear functional equations to the case of ϕ -pseudo-monotone mappings—a new and rather general class of mappings. An application is considered. © 1994 Academic Press, Inc.

1. INTRODUCTION

Petryshyn [2] considered an approximation scheme and applied it to the general approximation-solvability of nonlinear equations in general normed spaces. This scheme is specifically more application-oriented toward the numerical methods in the sense that solutions of an equation in some function space X can be approximated by solutions of equations in some discrete space, say $X_n = \mathbf{R}^n$. As an example, we can consider an integral equation

$$x(t) = \int_0^1 k(t, s)x(s) ds + f(t) \quad (1.1)$$

on $C([0, 1])$ with $k \in C([0, 1] \times [0, 1])$, and apply the convergent mechanical quadrature to approximate $\int_0^1 y(s) ds$, that is, we write

$$\int_0^1 y(s) ds = \sum_{j=1}^n a_{nj}y(s_{nj}) + r_n(y) \quad (1.2)$$

with coefficients $a_{nj} \geq 0$, interpolation points $0 \leq s_{n1} < \cdots < s_{nn} \leq 1$ and the remainder r_n such that $r_n(y) \rightarrow 0$ on $C([0, 1])$. Thus, the integral

equation is approximated by the discrete system

$$c_{ni} = \sum_{j=1}^n a_{nj} k(s_{ni}, s_{nj}) c_{nj} + f(s_{ni}) \quad \text{for } i = 1, \dots, n, \quad (1.3)$$

obtained by neglecting r_n and taking $t = s_{ni}$ for $i = 1, \dots, n$.

The purpose of this paper is twofold. First, we generalize the results on approximation-solvability, by applying the theorem of Petryshyn [2] to the case of ϕ -pseudo-monotone mappings—a new class of mappings in the sense of Vainberg [4]. Second, we give an application to the numerical range—a generalization of the Zarantonello numerical range [7].

DEFINITION 1.1. Let X and Y be complex Banach spaces, and $\phi: X \rightarrow Y$ be such that

- (i) $\phi(X)$ is dense in Y , and
- (ii) for each $x \in X$ and each $t \geq 0$, $\phi(tx) = t\phi(x)$.

A mapping $A: X \rightarrow Y^*$ from X to Y^* , the dual of Y , is said to be ϕ -pseudo-monotone if there exists a constant $d > 0$ such that

$$|[Ax - Au, \phi(x - u)]| \geq d \|x - u\| \|\phi(x - u)\| \quad \text{for all } x, u \in X, \quad (1.4)$$

where $[\cdot, \cdot]$ is the pairing between Y^* and Y .

Consequently,

$$|[Ax - Au, \phi(x - u)]| \geq d \|x\| - \|u\| \|\phi(x - u)\| \quad \text{for all } x, u \in X. \quad (1.5)$$

A is pseudo-monotone in the sense of Vainberg [4] when $Y = X$ (reflexive) and $\phi = I$.

However, usually pseudo-monotonicity carries a different meaning [8].

2. APPROXIMATION-SOLVABILITY

Suppose X is a separable reflexive complex Banach space with $\dim X = \infty$, and $A: X \rightarrow X^*$ from X to its dual X^* is ϕ -pseudo-monotone and continuous. Let (X_n) be a Galerkin scheme in X such that

$$X_n = \text{span}(e_{1n}, \dots, e_{n'n}), \quad n = 1, 2, \dots \quad (2.1)$$

Let $E_n: X_n \rightarrow X$ be the embedding mapping corresponding to $X_n \subseteq X$. We construct $R_n: X \rightarrow X_n$ in such a manner that, for each $x \in X$, there exists at

least an element $R_n x$ such that

$$\|x - R_n x\| = \text{dist}(x, X_n). \quad (2.2)$$

We consider the equation

$$Ax = b, \quad x \in X, b \in X^*, \quad (2.3)$$

along with approximate equations

$$E_n^* A E_n x_n = E_n^* b, \quad x_n \in X_n, n = 1, 2, \dots, \quad (2.4)$$

with respect to an approximation scheme $\pi = \{X_n, \phi_n, E_n, R_n, X_n^*, E_n^*\}$, represented by a diagram

$$\begin{array}{ccccccc} X & \xrightarrow{\phi} & X & \xrightarrow{A} & X^* & & \\ E_n \uparrow & & R_n \downarrow & E_n & & \downarrow E_n^* & \\ X_n & \xrightarrow{\phi_n} & X_n & \xrightarrow{A_n} & X_n^* & & \end{array} \quad (2.5)$$

where $A_n = E_n^* A E_n$ and $\phi_n = R_n \phi E_n$ are continuous. For more details on approximation-solvability, see Deimling [1], Petryshyn [2], Rosen [3], and Zeidler [8].

For $n = 1, 2, \dots$, approximate equations (2.4) are equivalent to Galerkin equations

$$[Ax_n, e_{jn}] = [b, e_{jn}] \quad (2.6)$$

for $x_n \in X_n, j = 1, \dots, n'$.

Here and in what follows, operators $E_n: X_n \rightarrow X$ and $E_n^*: X^* \rightarrow X_n^*$ are linear and continuous. The symbols " \rightarrow " and " \xrightarrow{w} " shall denote the strong convergence and weak convergence, respectively. Also, \mathbf{K} denotes the real or complex field.

We recall some definitions for the sake of the completeness.

DEFINITION 2.1 (Compatibility). The approximation scheme $\pi = \{X_n, \phi_n, E_n, R_n, X_n^*, E_n^*\}$ is called *compatible* if

$$\lim_{n \rightarrow \infty} \|E_n R_n x - x\|_X = 0 \quad \text{for all } x \in X. \quad (2.7)$$

DEFINITION 2.2 (Admissible Inner Approximation Scheme). The approximation scheme π is said to be an *admissible inner approximation scheme* iff

- (A1) X and X^* are infinite-dimensional normed spaces over \mathbf{K} .
 (A2) X_n and X_n^* are normed spaces over \mathbf{K} with $\dim X_n = \dim X_n^* < \infty$ for all n .
 (A3) $E_n: X_n \rightarrow X$ and $E_n^*: X^* \rightarrow X_n^*$ are linear and continuous with $\sup \|E_n\| < \infty$ and $\sup \|E_n^*\| < \infty$.
 (A4) The compatibility condition is satisfied.

DEFINITION 2.3 (Solvability). The equation $Ax = b$ is said to be *solvable* if it has a solution for each $b \in X^*$.

DEFINITION 2.4 (Unique Approximation-Solvability). The equation $Ax = b$ is called uniquely *approximation-solvable* if, for each $b \in X^*$, the following hold:

- (i) $Ax = b$, $x \in X$, has a unique solution.
 (ii) For each $n \geq n_0$, the approximate equation

$$E_n^* A E_n x_n = E_n^* b, \quad x_n \in X_n, \quad (2.8)$$

has a unique solution.

- (iii) The sequence (x_n) converges to the solution x of the equation $Ax = b$ in the sense that

$$\lim_{n \rightarrow \infty} \|E_n x_n - x\|_X = 0. \quad (2.9)$$

DEFINITION 2.5 (A-Properness). The operator $A: X \rightarrow X^*$ is *approximation-proper* (abbreviated as *A-proper*) with respect to the approximation scheme π if the following holds:

Let (n') be any subsequence of the sequence of natural numbers. If $(x_{n'})$ is a sequence with $x_{n'} \in X_{n'}$ for all n' , and if

$$\lim_{n' \rightarrow \infty} \|A_{n'} x_{n'} - E_{n'}^* b\|_{X_{n'}^*} = 0 \text{ for fixed } b \in X^* \quad \text{and} \quad \sup \|x_{n'}\| < \infty, \quad (2.10a), (2.10b)$$

then there exists a subsequence $(x_{n''})$ such that

$$\lim_{n'' \rightarrow \infty} \|E_{n''} x_{n''} - x\|_X = 0 \quad \text{and} \quad Ax = b. \quad (2.11a), (2.11b)$$

DEFINITION 2.6 (Consistency). The approximation scheme π is called *consistent* if, for all $x \in X$,

$$\lim_{n \rightarrow \infty} \|E_n^* Ax - A_n R_n x\|_{X_n^*} = 0. \quad (2.12)$$

DEFINITION 2.7 (Stability). The approximation scheme π is said to be *stable* if there exists an n_0 such that

$$\|A_n x - A_n u\|_{X_n^*} \geq d \|x - u\|_{X_n} \quad (2.13)$$

for all $x, u \in X_n$ and $n \geq n_0$, where $d > 0$.

Before proceeding to the main result, we need to state the crucial theorem of Petryshyn [2] on the inner approximation-solvability.

LEMMA 2.1. *If the approximation scheme $\pi = \{X_n, \phi_n, E_n, R_n, X_n^*, E_n^*\}$ is an admissible inner approximation scheme with consistency and stability, then the equation*

$$Ax = b, \quad x \in X, \quad (2.14)$$

is uniquely approximation-solvable for each $b \in X^$ iff A is A -proper.*

Now we establish the main result on the approximation-solvability of nonlinear equations involving ϕ -pseudo-monotone operators.

THEOREM 2.1. *Let X be a separable reflexive complex Banach space with $\dim X = \infty$. Let $\pi = \{X_n, \phi_n, E_n, R_n, X_n^*, E_n^*\}$ be an approximation scheme for the pair (X, X^*) represented by the diagram (2.5), and let $\phi: X \rightarrow X$ be continuous such that*

$$R_n \phi E_n x = \phi x \quad \text{for all } x \in X_n \text{ and each } n. \quad (2.15)$$

If $A: X \rightarrow X^$ is ϕ -pseudo-monotone and continuous, then, for each $b \in X^*$, the equation*

$$Ax = b, \quad x \in X, \quad (2.16)$$

is uniquely approximation-solvable.

For $\phi = I$, we arrive at the following.

COROLLARY 2.1. *If X is a separable reflexive complex Banach space with $\dim X = \infty$, $\pi_0 = \{X_n, E_n, R_n, X_n^*, E_n^*\}$ is an approximation scheme for the pair (X, X^*) , and $A: X \rightarrow X^*$ is pseudo-monotone and continuous, then, for each $b \in X^*$, the equation*

$$Ax = b, \quad x \in X, \quad (2.17)$$

is uniquely approximation-solvable.

Proof of Theorem 2.1. As our method of proof is based upon Lemma 2.1, we first show that the approximation scheme $\pi = \{X_n, \phi_n, E_n, R_n, X_n^*,$

E_n^* is an admissible inner approximation scheme such that π is consistent and stable. Second, we show that $A: X \rightarrow X^*$ is A -proper.

Admissible Inner Approximation Scheme. Since $\|E_n\| = 1$ (and hence $\|E_n^*\| = 1$) and (X_n) is a Galerkin scheme, $\text{dist}(x, X_n) \rightarrow 0$ as $n \rightarrow \infty$ for all $x \in X$. This implies that $\|R_n x - x\| \rightarrow 0$ as $n \rightarrow \infty$, and thus, the compatibility condition is satisfied.

Consistency. Since A is continuous and the compatibility condition (2.7) is satisfied, the consistency condition is as follows: Since

$$\|AE_n R_n x - Ax\| \rightarrow 0 \quad \text{as } n \rightarrow \infty \text{ and } \|E_n^*\| = 1, \quad (2.18)$$

this implies that, as $n \rightarrow \infty$,

$$\|E_n^* Ax - A_n R_n x\| = \|E_n^* Ax - E_n^* AE_n R_n x\| \leq \|E_n^*\| \|Ax - AE_n R_n x\| \rightarrow 0. \quad (2.19)$$

Stability. For all $x, y \in X_n$, we obtain

$$\begin{aligned} \|A_n x - A_n y\| \|\phi(x - y)\| &\geq |[A_n x - A_n y, \phi(x - y)]| \\ &= |[E_n^* AE_n x - E_n^* AE_n y, \phi(x - y)]| \\ &= |[AE_n x - AE_n y, E_n \phi(x - y)]| \\ &= |[Ax - Ay, \phi(x - y)]| \\ &\geq d\|x - y\| \|\phi(x - y)\|, \end{aligned} \quad (2.20)$$

and consequently,

$$\|A_n x - A_n y\| \geq d\|x - y\| \quad \text{for all } x, y \in X_n. \quad (2.21)$$

A-Properness. Assume $\sup\|x_n\| < \infty$ for all $x_n \in X_n$, and

$$\|A_n x_n - E_n^* b\| = \|E_n^* Ax_n - E_n^* b\| \rightarrow 0 \quad \text{as } n \rightarrow \infty \text{ for all } n. \quad (2.22)$$

Since X is reflexive, there exists a subsequence, again denoted by (x_n) , such that

$$x_n \xrightarrow{w} x \quad \text{in } X \text{ as } n \rightarrow \infty. \quad (2.23)$$

This means that it suffices to show that

$$x_n \rightarrow x \text{ as } n \rightarrow \infty \quad \text{and} \quad Ax = b. \quad (2.24)$$

Since $\|R_n x - x\| \rightarrow 0$ as $n \rightarrow \infty$, $x_n \xrightarrow{w} x$ as $n \rightarrow \infty$ implies that

$$x_n - R_n x \xrightarrow{w'} 0 \quad \text{as } n \rightarrow \infty. \quad (2.25)$$

Thus, as $n \rightarrow \infty$, with (2.22) we get

$$\begin{aligned} \|E_n^* A x_n - E_n^* A R_n x\| &= \|(E_n^* A x_n - E_n^* b) + (E_n^* b - E_n^* A R_n x)\| \\ &\leq \|E_n^* b - E_n^* A x\| + \|E_n^* (b - A x)\|. \end{aligned} \quad (2.26)$$

It follows from condition (2.20) and the above results that we get, for $x_n \in X_n$ as above, as $n \rightarrow \infty$,

$$\begin{aligned} d\|x_n - R_n x\| \|\phi(x_n - R_n x)\| &\leq |A_n x_n - A_n R_n x, \phi(x_n - R_n x)| \\ &= |[E_n^* A x_n - E_n^* A R_n x, \phi(x_n - R_n x)]| \rightarrow 0. \end{aligned} \quad (2.27)$$

This also implies that $x_n - R_n x \rightarrow 0$ as $n \rightarrow \infty$. Since $\sup\|E_n\| < \infty$, we find

$$\|E_n x_n - E_n R_n x\| \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (2.28)$$

and the compatibility condition gives

$$\|E_n R_n x - x\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (2.29)$$

Thus,

$$\|E_n x_n - x\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (2.30)$$

It only remains to show that $Ax = b$. Since $x_n - R_n x \rightarrow 0$ as $n \rightarrow \infty$ implies that $x_n \rightarrow x$ as $n \rightarrow \infty$, and since A is continuous, we obtain $Ax = b$. Hence A is A -proper.

Now the equation $Ax = b$ is uniquely approximation-solvable by Lemma 2.1, and this completes the proof.

3. APPLICATION TO NUMERICAL RANGE

This section deals with the approximation-solvability of the equations involving numerical ranges.

DEFINITION 3.1 (Numerical Range). Let $A: X \rightarrow X^*$ be a mapping from a separable reflexive complex Banach space X to its dual X^* . The set

$$n_0[A] = \left\{ \frac{[Ax, \phi x] + [Ay - Az, \phi(y - z)]}{[Jx, \phi x] + [Jy - Jz, \phi(y - z)]} : x, y, z \in X, y \neq z \right\} \quad (3.1)$$

is called the *numerical range* of A , where $J: X \rightarrow X^*$ is strictly monotone normalized duality, and $\phi: X \rightarrow X$ is continuous such that

- (i) $\phi(X)$ is dense in X ; and
- (ii) for each $t \geq 0$, $\phi(tx) = t\phi(x)$.

A continuous function $a: R^+ \rightarrow R^+$ is called a *gauge function* if $a(0) = 0$, and a is strictly increasing. Let X be a real reflexive Banach space and X^* its dual. A mapping $J: X \rightarrow X^*$ is called a *duality mapping* between X and X^* with respect to a gauge function a if

$$[Jxx] = a(\|x\|)\|x\| \text{ and } \|Jx\| = a(\|x\|) \quad \text{for } x \in X. \quad (3.2)$$

We note that if $a(t) = t$, J is said to be a *normalized duality mapping*. If X^* is strictly convex, then J is uniquely determined by a , and if X is also reflexive, then J is a single-valued semicontinuous mapping of X onto X^* , which is bounded and positively homogenous. Furthermore, J is monotone and satisfies, for all $x, y \in X$,

$$[Jx - Jy, x - y] \geq (a(\|x\|) - a(\|y\|))(\|x\| - \|y\|). \quad (3.3)$$

Also, we have

$$[Jx - Jy, x - y] = [Jx, x - y] - [Jy, x - y] \geq |a(\|x\|) - a(\|y\|)| \|x - y\|. \quad (3.4)$$

If J is normalized duality, then

$$[Jx - Jy, x - y] \geq (\|x\| - \|y\|)^2 \quad \text{for all } x, y \in X, \quad (3.5)$$

which implies that

$$[Jx - Jy, x - y] \geq |\|x\| - \|y\|| \|x - y\| \quad \text{for all } x, y \in X. \quad (3.6)$$

Next, we describe some of the properties of the numerical range $n_0[A]$ along with some special cases of significance.

Clearly, $n_0[A]$ reduces to the numerical range (Verma [6]) defined by

$$V[A] = \left\{ \frac{\langle Ax, x \rangle + \langle Ay - Az, y - z \rangle}{\|x\|^2 + \|y - z\|^2} : x, y, z \in X, y \neq z \right\}, \quad (3.7)$$

where X is a Hilbert space and ϕ is the identity. Furthermore, $n_0[A]$ coincides with the Zarantonello numerical range [7] defined by

$$\mathcal{N}[A] = \left\{ \frac{\langle Ay - Az, y - z \rangle}{\|y - z\|^2} : y, z \in X, y \neq z \right\} \quad (3.8)$$

where X is a Hilbert space with $\phi = I$ and $x = 0$.

THEOREM 3.1. *Let $A, B: X \rightarrow X^*$ be mappings from a reflexive Banach space X to its dual X^* , and $\lambda \in \mathbf{K}$. Then*

- (i) $n_0[\lambda A] = \lambda n_0[A]$.
- (ii) $n_0[A + B] \subset n_0[A] + n_0[B]$.
- (iii) $n_0[A - \lambda J] = n_0[A] - \{\lambda\}$,

where $J: X \rightarrow X^*$ is strictly monotone normalized duality.

Proof. The proof follows from the Definition 3.1.

We now establish the main result of the section.

THEOREM 3.2. *Let $A: X \rightarrow X^*$ be continuous from a separable reflexive complex Banach space X to its dual X^* . If X and X^* are locally uniformly convex, a number $\lambda \in \mathbf{K}$ has a positive distance from the numerical range of A , $n_0[A]$, that is,*

$$d = \inf\{|\lambda - \mu| : \mu \in n_0[A]\} > 0, \quad (3.9)$$

$\phi: X \rightarrow X$ is continuous, and $J: X \rightarrow X^$ is a normalized duality, then the equation*

$$Ax - \lambda Jx = b, \quad x \in X, \quad (3.10)$$

is uniquely approximation-solvable for each $b \in X^$.*

COROLLARY 3.1. *Theorem 3.2 reduces to a result of the author [6, Theorem 2.2] when X is a Hilbert space and ϕ is the identity.*

We derive the following corollary when X is a Hilbert space with $x = 0$ and $\phi = I$.

COROLLARY 3.2. (Zeidler [8, Theorem 34.C]). *Suppose $A: X \rightarrow X$ is continuous from a separable Hilbert space X into itself. If a number $\lambda \in \mathbf{K}$ has a positive distance from the Zarantonello numerical range of A , $\mathcal{N}[A]$, that is,*

$$d = \inf\{|\lambda - \mu| : \mu \in \mathcal{N}[A]\} > 0, \quad (3.11)$$

then the equation

$$Ax - \lambda Ix = b, \quad x \in X, \quad (3.12)$$

is uniquely approximation-solvable for each $b \in X$.

Proof of Theorem 3.2. Before we can apply Theorem 2.1, we need to obtain the key estimates. Since $J: X \rightarrow X^*$ is strictly monotone, we find, for $x, y, z, \in X$ with $y \neq z$,

$$\begin{aligned} & |[(A - \lambda J)x, \phi x] + [(A - \lambda J)y - (A - \lambda J)z, \phi(y - z)]| \\ &= |[Ax, \phi x] - \lambda[Jx, \phi x] + [Ay - Az, \phi(y - z)] \\ &\quad - \lambda[Jy - Jz, \phi(y - z)]| \\ &= |[Ax, \phi x] + [Ay - Az, \phi(y - z)] - \lambda([Jx, \phi x] \\ &\quad + [Jy - Jz, \phi(y - z)])| \quad (3.13) \\ &= \left| \frac{[Ax, \phi x] + [Ay - Az, \phi(y - z)]}{[Jx, \phi x] + [Jy - Jz, \phi(y - z)]} - \lambda \right| |[Jx, \phi x] \\ &\quad + [Jy - Jz, \phi(y - z)]| \\ &\geq d \operatorname{Re}([Jx, \phi x] + [Jy - Jz, \phi(y - z)]) \\ &\geq d(\|x\| \|\phi x\| + \|y\| - \|z\| \|\phi(y - z)\|). \end{aligned}$$

For $x = 0$, we obtain an estimate

$$|[A - \lambda J)y - (A - \lambda J)z, \phi(y - z)]| \geq d\|\|y\| - \|z\|\|\phi(y - z)\| \quad (3.14)$$

for all $y, z \in X$.

The proof can be obtained from that of Theorem 2.1.

ACKNOWLEDGMENT

This work was supported by a research grant to the second author from the University of Central Florida. The first author was a research associate of this project. The authors express their grateful thanks to the referee for suggesting some improvements.

REFERENCES

1. K. DEIMLING, "Nonlinear Functional Analysis," Springer-Verlag, New York, 1985.
2. W. PETRYSHYN, Nonlinear equations involving noncompact operators, in "Nonlinear Functional Analysis," Proc. Symposia in Pure Math., Chicago, 1968, Vol. 18, pp. 206-233, Amer. Math. Soc., Providence, RI, 1970.

3. I. G. ROSEN, Convergence of Galerkin approximation for operators Riccati equations—A nonlinear evolution equation approach, *J. Math. Anal. Appl.* **155** (1991), 226–248.
4. M. M. VAINBERG, "Variational Method and Method for Monotone Operators in the Theory of Nonlinear Equations," Wiley, New York, 1973.
5. R. VERMA, Joint spatial numerical range of tensor products, *Proc. Roy. Irish Acad. A* **90** (1990), 201–204.
6. R. VERMA, Role of numerical range in approximation-solvability of nonlinear functional equations, *Appl. Math. Lett.* **5** No. 2 (1992), 25–27.
7. E. ZARANTONELLO, The closure of the numerical range contains the spectrum, *Bull. Amer. Math. Soc.* **70** (1964), 781–787.
8. E. ZEIDLER, "Nonlinear Functional Analysis and Its Applications II/B," Springer-Verlag, New York, 1990.